

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Acc65I
Catalog Number: R0599S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBC4 DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number:10044861Expiration Date:05/2021Storage Temperature:-20°C

Storage Conditions: 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0599S/L v1.0

Acc65I Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0599SVIAL	Acc65I	10044862	Pass	
B7203SVIAL	NEBuffer™ 3.1	10041637	Pass	

Assay Name/Specification	Lot # 10044861
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 50 Units of Acc65l incubated for 4 hours at 37°C results in <20%	Pass
conversion to the nicked form as determined by agarose gel electrophoresis. Blue-White Screening (Terminal Integrity) A sample of Litmus28i vector linearized with a 10-fold excess of Acc65I, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene	Pass
results in <1% white colonies. Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of Acc65l incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of pBC4 DNA and a minimum of 100 Units of Acc65l incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass



R0599S / Lot: 10044861

Page 1 of 2



Assay Name/Specification	Lot # 10044861
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pBC4 DNA with Acc65I, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Acc65I.	Pass

This product has been tested and shown to be in compliance with all specifications.

Production Scientist 05 Apr 2019

Michael Tonello Packaging Quality Co

Packaging Quality Control Inspector 03 Jul 2019



R0599S / Lot: 10044861

Page 2 of 2