

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Acc65I
Catalog Number: R0599S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBC4 DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10172236
Expiration Date: 11/2024
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0599S/L v1.0

Acc65I Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0599SVIAL	Acc65I	10172235	Pass	
B6003SVIAL	NEBuffer™ r3.1	10168652	Pass	

Assay Name/Specification	Lot # 10172236
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of pBC4 DNA and a minimum of 100	Pass
Units of Acc65I incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pBC4 DNA with Acc65I, >95% of the DNA fragments	Pass
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with Acc65I.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 μl reaction in NEBuffer 3.1 containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of Acc65l incubated for	
4 hours at 37°C releases <0.1% of the total radioactivity.	
Endonuclease Activity (Nicking)	Pass
A 50 μl reaction in NEBuffer 3.1 containing 1 μg of supercoiled PhiX174 DNA and a	
minimum of 50 Units of Acc65I incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	



R0599S / Lot: 10172236

Page 1 of 2

Assay Name/Specification	Lot # 10172236
Blue-White Screening (Terminal Integrity) A sample of Litmus28i vector linearized with a 10-fold excess of Acc65I, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Suń

Production Scientist

24 Nov 2022

Michael Tonello

Packaging Quality Control Inspector

28 Nov 2022



R0599S / Lot: 10172236

Page 2 of 2