

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Nspl
Catalog Number: R0602L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10173539
Expiration Date: 12/2024
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 0.15% Triton X-100, 200 µg/ml BSA

Specification Version: PS-R0602S/L v1.0

Nspl Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0602LVIAL	Nspl	10173538	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10175289	Pass	
B6004SVIAL	rCutSmart™ Buffer	10173160	Pass	

Assay Name/Specification	Lot # 10173539
Protein Purity Assay (SDS-PAGE) NspI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue	Pass
detection.	
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of supercoiled PhiX174 DNA and	Pass
a minimum of 10 units of NspI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of Nspl incubated for 4	
hours at 37°C releases <0.1% of the total radioactivity.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of Lambda DNA with NspI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with Nspl.	



R0602L / Lot: 10173539

Page 1 of 2

Assay Name/Specification	Lot # 10173539
Non-Specific DNase Activity (16 Hour)	Pass
A 50 μI reaction in CutSmart™ Buffer containing 1 μg of Lambda DNA and a minimum of	
50 Units of NspI incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun

Production Scientist 09 Jan 2023 Michael Tonello

Packaging Quality Control Inspector

10 Jan 2023



R0602L / Lot: 10173539

Page 2 of 2