240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Nt.BstNBI

Catalog #: R0607S/L

Concentration: 10,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg T7 DNA in 1 hour at 55°C in a total reaction volume of

 $50 \,\mu l$.

 Lot #:
 0401605

 Assay Date:
 05/2016

 Expiration Date:
 05/2018

 Storage Temp:
 -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA

Specification Version: PS-R0607S/L v1.0
Effective Date: 26 Oct 2015

Assay Name/Specification (minimum release criteria)	Lot #0401605
Exonuclease Activity (Radioactivity Release) - A 50 μ l reaction in NEBuffer 3.1 containing 1 μ g of a mixture of single and double-stranded [3 H] <i>E. coli</i> DNA and a minimum of 50 units of Nt.BstNBI incubated for 4 hours at 55°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of T7 DNA with Nt.BstNBI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Nt.BstNBI.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of T7 DNA and a minimum of 10 Units of Nt.BstNBI incubated for 16 hours at 55°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) - Nt.BstNBI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass

Authorized by Derek Robinson 26 Oct 2015







Inspected by Anthony Francis 06 May 2016