

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Btgl
Catalog Number: R0608L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBR322 DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number: 10008618
Expiration Date: 04/2020
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 μg/ml BSA

Specification Version: PS-R0608S/L v1.0

| Btgl Component List    |                       |            |                      |  |
|------------------------|-----------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description | Lot Number | Individual QC Result |  |
| R0608LVIAL             | Btgl                  | 0071804    | Pass                 |  |
| B7204SVIAL             | CutSmart® Buffer      | 3091805    | Pass                 |  |

| Assay Name/Specification   | Lot # 10008618 |
|--|----------------|
| Protein Purity Assay (SDS-PAGE)  | Pass           |
| Btgl is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.  |                |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 50 units of Btgl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.                | Pass           |
| Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pBR322 DNA with Btgl, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Btgl.  | Pass           |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of 50 Units of Btgl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |

This product has been tested and shown to be in compliance with all specifications.



R0608L / Lot: 10008618 Page 1 of 2



Penghua Zhang Production Scientist 27 Jun 2018 Michael Tonello

Packaging Quality Control Inspector

27 Jun 2018

