

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	BtsI-v2
Catalog Number:	R0667S
Concentration:	10,000 U/mI
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μ l.
Packaging Lot Number:	10122954
Expiration Date:	07/2023
Storage Temperature:	-20°C
Storage Conditions:	50 mM KCl , 10 mM Tris-HCl, 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 200 μg/ml BSA (pH 7.4 @ 25°C)
Specification Version:	PS-R0667S/L v3.0

BtsI-v2 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0667SVIAL	BtsI-v2	10114800	Pass	
B6004SVIAL	rCutSmart™ Buffer	10123108	Pass	

Assay Name/Specification	Lot # 10122954
Protein Purity Assay (SDS-PAGE) BtsI-v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with BtsI-v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with BtsI-v2.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 10 units of BtsI-v2 incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of BtsI-v2 incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10122954
Functional Testing (15 minute Digest)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and 1 µl of	
BtsI-v2 incubated for 15 minutes at 37°C results in complete digestion as determined	
by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Panatha Thang

Penghua Zhang Production Scientist 20 Oct 2021

Michae

Michael Tonello Packaging Quality Control Inspector 20 Oct 2021

