

## New England Biolabs Certificate of Analysis

**Product Name:** BspQI  
**Catalog #:** R0712S/L  
**Concentration:** 10,000 units/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 50°C in a total reaction volume of 50 µl.  
**Lot #:** 0021706  
**Assay Date:** 06/2017  
**Expiration Date:** 6/2019  
**Storage Temp:** -20°C  
**Storage Conditions:** 500 mM KCl , 20 mM Tris-HCl (pH 7.0), 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 0.10 % TritonX-100 , 500 µg/ml BSA  
**Specification Version:** PS-R0712S/L v2.0  
**Effective Date:** 30 Oct 2015

Assay Name/Specification (minimum release criteria)	Lot #0021706
<b>Endonuclease Activity (Nicking)</b> - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled M13mp18 DNA and a minimum of 10 units of BspQI incubated for 4 hours at 50°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 50 units of BspQI incubated for 4 hours at 50°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of Lambda DNA with BspQI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BspQI.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 hour)</b> - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 10 Units of BspQI incubated for 16 hours at 50°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.	<b>Pass</b>
<b>Protein Purity Assay (SDS-PAGE)</b> - BspQI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	<b>Pass</b>

---

## New England Biolabs Certificate of Analysis

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



---

Authorized by  
Derek Robinson  
30 Oct 2015



---

Inspected by  
Penghua Zhang  
02 Jun 2017

