

New England Biolabs Certificate of Analysis

Product Name: *EagI-HF®*
Catalog Number: *R3505M*
Concentration: *100,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 µl.*
Packaging Lot Number: *10121634*
Expiration Date: *09/2023*
Storage Temperature: *-80°C*
Storage Conditions: *500 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 200 µg/ml BSA, (pH 7.4 @ 25°C)*
Specification Version: *PS-R3505M v3.0*

EagI-HF® Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R3505MVIAl	EagI-HF®	10121633	Pass
B7024AVIAl	Gel Loading Dye, Purple (6X)	10108733	Pass
B6004SVIAl	rCutSmart™ Buffer	10120518	Pass

Assay Name/Specification	Lot # 10121634
Protein Purity Assay (SDS-PAGE) EagI-HF™ is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of EagI-HF™ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Blue-White Screening (Terminal Integrity) A sample of Litmus38i vector linearized with a 10-fold excess of EagI-HF™, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pXba DNA with EagI-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with EagI-HF™.	Pass

Assay Name/Specification	Lot # 10121634
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 100 Units of EagI-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 Units of EagI-HF™ incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Penghua Zhang
Production Scientist
01 Oct 2021



Michael Tonello
Packaging Quality Control Inspector
01 Oct 2021