

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Agel-HF®
Catalog Number: R3552L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total

reaction of 50 μl.

Packaging Lot Number: 10160219
Expiration Date: 04/2024
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200

μg/ml rAlbumin (pH 7.4 @ 25°C)

Specification Version: PS-R3552S/L v2.0

Agel-HF® Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R3552LVIAL	Agel-HF®	10145782	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10161525	Pass	
B6004SVIAL	rCutSmart™ Buffer	10156434	Pass	

Assay Name/Specification	Lot # 10160219
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of	
100 units of Agel-HF® incubated for 16 hours at 37°C results in a DNA pattern free	
of detectable nuclease degradation as determined by agarose gel electrophoresis.	
igation and Recutting (Terminal Integrity)	Pass
After a 20-fold over-digestion of Lambda DNA with Agel-HF®, >95% of the DNA	
ragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	
fragments, >95% can be recut with Agel-HF®.	
Functional Testing (15 minute Digest)	Pass
λ 50 μl reaction in rCutSmart™ Buffer containing 1 μg of Lambda DNA and 1 μl of	
Agel-HF® incubated for 15 minutes at 37°C results in complete digestion as	
determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of a mixture of single and	



R3552L / Lot: 10160219

Page 1 of 2

Assay Name/Specification	Lot # 10160219
double-stranded [ ³H] E. coli DNA and a minimum of 200 units of Agel-HF® incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Blue-White Screening (Terminal Integrity) A sample of LITMUS28i vector linearized with a 10-fold excess of Agel-HF®, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
<b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 20 units of Agel-HF® is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
Protein Purity Assay (SDS-PAGE)  Agel-HF® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

07 Sep 2022

Erin Varney

Packaging Quality Control Inspector

07 Sep 2022



R3552L / Lot: 10160219

Page 2 of 2