

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Apol-HF®
Catalog Number: R3566S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37 degrees C in a total reaction volume

of 50 μL

Packaging Lot Number: 10126513
Expiration Date: 08/2023
Storage Temperature: -20°C

Storage Conditions: 200 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol,

200 $\mu g/ml$ BSA, (pH 7.4 @ 25°C)

Specification Version: PS-R3566S/L v1.0

Apol-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3566SVIAL	Apol-HF®	10116139	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10121393	Pass	
B6004SVIAL	rCutSmart™ Buffer	10124441	Pass	

Assay Name/Specification	Lot # 10126513
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and	Pass
double-stranded [³H] E. coli DNA and a minimum of 100 units of Apol-HF incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with Apol-HF, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Apol-HF.	Pass
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of Apol-HF, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of	Pass



R3566S / Lot: 10126513

Page 1 of 2

Assay Name/Specification	Lot # 10126513
100 units of Apol-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) Apol-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Functional Testing (15 minute Digest) A 50 μl reaction in CutSmart® Buffer containing 1 μg of Lambda DNA and 1 μl of Apol-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

29 Nov 2021

Michael Tonello

Packaging Quality Control Inspector

29 Nov 2021



R3566S / Lot: 10126513

Page 2 of 2