

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: SbfI-HF®
Catalog Number: R3642L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10083545
Expiration Date: 02/2022
Storage Temperature: -20°C

Storage Conditions: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R3642S/L v1.0

SbfI-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3642LVIAL	SbfI-HF®	10067430	Pass	
B7204SVIAL	CutSmart® Buffer	10081171	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10084970	Pass	

Assay Name/Specification	Lot # 10083545
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 20 Units of Sbfl-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of	Pass
detectable nuclease degradation as determined by agarose gel electrophoresis. Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of Lambda DNA with SbfI-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SbfI-HF™.	
Protein Purity Assay (SDS-PAGE) SbfI-HF™ is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of Sbfl-HF™ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass



R3642L / Lot: 10083545

Page 1 of 2

Assay Name/Specification	Lot # 10083545
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 20 Units of Sbfl-HF™ incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang **Production Scientist**

15 Oct 2020

Michael Tonello

Packaging Quality Control Inspector

15 Oct 2020

