

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name:	Thermolabile Proteinase K
Catalog #:	P8111S
Concentration:	120 units/ml
Unit Definition:	One unit is defined as the amount of enzyme required to release 1.0 μ mol of 4-nitroaniline per minute from N-Succinyl-Ala- Ala-Pro-Phe-p-nitroanilide at 25°C in a total reaction volume of 105 μ l.
Shelf Life:	24 months
Storage Temp:	-20°C
Storage Conditions:	20 mM Tris-HCl, 1 mM CaCl ₂ , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version:	PS-P8111S v1.0
Effective Date:	19 Oct 2018

Assay Name/Specification (minimum release criteria)

qPCR DNA Contamination (Eukaryotic Genomic) - A minimum of 0.12 units of Thermolabile Proteinase K is screened for the presence of eukaryotic genomic DNA using SYBR® Green qPCR with universal primers for the 18S rRNA locus. Results are quantified using a standard curve generated from purified *E. album* genomic DNA. The measured level of eukaryotic genomic DNA contamination is $\leq 2.5 \text{ pg DNA/µl}$.

qPCR DNA Contamination (E. coli Genomic) - A minimum of 0.12 units of Thermolabile Proteinase K is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.

RNase Activity (Extended Digestion) - A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ l of Thermolabile Proteinase K is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

Single Stranded DNase Activity (FAM-Labeled Oligo) - A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 0.6 units of Thermolabile Proteinase K incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.

Date 19 Oct 2018

Derek Robinson Director of Quality Control



PS-P8111S v1.0 Page 1 of 1